

EFFICIENCY }

OFFICE PAD
361



Notes on the

Nesting of Birds

1935

J.E.B. Campbell
Big Stone Gap

April 4, 1935

EUROPEAN STARLING

5

FRESH

15' in hole in
maple - the hole (1' deep)
was slanting & was in a slanting
limb (1.5' in diam.) On V.E.S. campsite
[COMPOSED OF STRAW, FEATHERS, GRASS & PAPER]

1.18 x .83 1.18 x .83 ~~1.18 x .83~~
1.19 x .82 1.12 x .83
1.21 x .83

April 21, 1935

LOUISIANA WATER THRUSH

5

FRESH

Under some over-
hanging mass
about 3' above bed of a creek on
the bank - in deep woods 3/4 mi. V.E.
[COMPOSED OF MUDDY LEAVES, GRASSES AND
LINED WITH FINE GRASSES.]

.70 x .59
.72 x .58
.72 x .60
.72 x .58
.71 x .57

May 8, 1935

TURKEY VULTURE

(see Block NB)

2

$\frac{4}{5}$

Under a large rock about 30 yds. above James R. - The nest was about 2' in diam. (eggs were on bare ground) The entrance was about 3' high & at nest the height was about 1.5'

2.83 x 1.83

2.90 x 1.82

May 10, 1935

OVENBIRD

6

FRESH

On the ground near a log in a small clump of weeds. [composed of leaves and heavy grasses with a lining of hair and fine grasses]

.74 x .55

.72 x .56

.75 x .56

.73 x .54

.75 x .55

.73 x .55

May 9, 1935

WHIP-POOR-WILL

2

FRESH

Near Dorchester

Va. A slt. depression on bare
oak leaves - no nesting materials
- in grove of hardwoods at
(50 yds inside) edge of forest -

1.09 x .79

1.05 x .77

May 13, 1935

MARYLAND YELLOW THROAT

5

✓

FRESH

In a small
clump of weeds about 5" above
the ground. Under V.E.S. bridge.

[COMPOSED OF GRASSES, LEAVES, WEED STEMS AND A
FEW RUSHES - LINED WITH FINE GRASSES AND HAIR.]

.66 x .52

.69 x .52

.65 x .50

.68 x .51

.67 x .51

Date - May 13, 1935

No.

Name - COWBIRD

No. in set - 1

Incubation - FRESH

Situation - In a Marl. Yellow Throats nest (taken same day)

Sizes - .86 x .63

Date - May 16, 1935

No.

Name - BLUE-GRAY GNATCATCHER

No. in set - 4

Incubation - 1/5

Situation - In a horizontal limb of a walnut - about 15' from ground & about 5' from trunk of tree. The limb was about 1" in diam. (J. N. V.E.S.)

Sizes

[COMPOSED OF VEGETABLE FIBERS FINE ROOTLETS SOFT PLANT DOWN COVERED WITH LICHENS & BEAUTIFULLY BOUND & CAMOUFLAGED WITH SILKEN FIBERS & SPIDER WEB]
.54 x .44 .55 x .43
.56 x .43 .56 x .44 DIAM: 2 1/4" Inside 1 1/4" Ht. 2 1/2"

Date - May 22, 1935

No.

Name - YELLOW WARBLER

No. in set - 4

Incubation - FRESH

Situation - In a Three-pronged fork of a young gum. About 15' from the ground. [3/4 m. V.E.S.]

[COMPOSED OF SOFT PLANT FIBERS, LIGHT GRAPE-VINE BARK AND LINED WITH COTTONY SUBSTANCES AND HORSEHAIR]

Sizes

.62 x .48 .62 x .49
.61 x .48 .61 x .49

Date - May 22, 1935 No.

Name - SUMMER TANAGER

No. in Set - 4

Incubation - FRESH

Location - On a horizontal limb of a small oak about 10' above a small country road. I.M. V.E.S.
[COMPOSED OF WEEDS, GRASSES, AND LINED WITH FINE BLEACHED GRASSES]

Sizes - .91x.63 .92x.63
.90x.60 .88x.63

Date - May 25, 1935 No.

Name - SCARLET TANAGER

No. in Set - 4

Incubation - FRESH

Location - On a horizontal limb of a poplar about 8' above a road.
[COMPOSED OF WEEDS AND GRASSES AND LINED WITH FINER GRASSES - NOT AS COMPACT OR AS CAREFULLY BUILT AS THE SUMMER TANAGER]

Sizes -

Date - JUNE 10, 1935 No.

Name - LEAST BITTERN

No. in Set - 3

Incubation - $\frac{1}{2}$

Location - Among a lot of cat tails which were bent over to form the nest. About 2' above the water in a marsh at Virginia Beach.

Sizes -

Date - JUNE 11, 1935 No.

Name - WOOD PEWEE

No. in set - 3

Incubation - $\frac{1}{3}$

Location - On the horizontal
of a locust about 15' from
ground & 4' from trunk.

Sizes -

Date - JULY 6, 1935 No.

Name - RUBY THROATED HUMMINGBIRD

No. in set - 1 (there were 2 - 1
(sadly) broke one)

Incubation - FRESH

Location - On a limb of a
beech - about 15'
from ground. In river near
home.

Sizes -

Handwritten text, likely bleed-through from the reverse side. Includes the date "22nd 9 1872" and the name "J. G. ...".


Handwritten text, likely bleed-through from the reverse side. Includes the date "22nd 11 1872" and the name "J. G. ...".


NO	NAME	NO IN SET	DATE	INC	NO.	SITUATION	VICINITY
1	BLACK BIRD						
	RED WINGED	4	MAY 12, 1933	F	498		
2	BLUE BIRD				766		
	EASTERN CATBIRD						
3	COMMON CHICADEE				704		
4	BLACK CAPPED CHUCK-WILLS WIDOW				735		
5	COWBIRD				416		
6	CHAT				495		
7	YELLOW BREAST CUCKOO				683		
8	YELLOW BILL DOVE				387		
9	MOORNING FLICKER				316		
10	YELLOW SHAFY SNATCHER				412		
11	BLUE GRAY GRACKLE				751		
12	PURPLE GROSBARK				511		
13	CARIBAL HAWK				593		
14	FISH HAWK				364		
15	RED TAIL JAY						
16	BLUE KILLDEER				477		
17	KINGFISHER				273		
18	BELTED MEADOWLARK				390		
19	MOCKINGBIRD				501		
20	ORIOLE				703		
21	BALTIMORE ORIOLE				507		
22	ORCHARD OVENBIRD				506		
23	OWL				674		
24	SCHREECH PEWEE				373		
25	WOOD HUMMINGB.				461		
26	RUBY THROATED BITTERN				438		
27	LEAST				761		
28	ROBIN				761		
29	AMERICAN SANDPIPER				263		
30	SPOTTED SANDPIPER				246		
31	SEMIPALM SPARROW				560		
32	CHIP. SPARROW						
	ENGLISH						

33	SPARROW	581
	SONG	
34	SPARROW	558
	WHITE THR.	
	SPARROW	
35	FIELD	563
	SPARLING	
36	EUROPEAN	493
	TANAGER	
37	SCARLET	608
	TANAGER	
38	SUMMER	610
	THRASHER	
39	BROWN	705
	THRUSH	
40	LA. WREN	676
	THRUSH	
41	WOOD	755
	TOWHEE	
42		587
	VIREO	
43	RED EYED	624
	VIREO	
44	WHITE EYED	631
	SWALLOW	
45	ROUGH WING	617
	VULTURE	
46	TURKEY	
	WARBLER	
47	PID. YE. THR.	681
	WARBLER	
48	YELLOW	652
	WOODPECKER	
49	RED HEAD	406
	WREN	
50	CAROLINA	718
	WREN	
51	HOUSE	721
	WREN	
52	WINTER	722
	WHIP-POOR	
53	WILL	417
	CROW	
54	COMMON	488
	QUAIL	
55		289
56	PHOEBE	456

GRACKLE

BOAT TAIL





NOTE IN A CORN (HILL) -
- TOOK FIVE QUAIL EGGS

NOTE IN A CORN (HILL) -
- TOOK FIVE QUAIL EGGS

NOTE IN A CORN (HILL) -
- TOOK FIVE QUAIL EGGS

NOTE IN A CORN (HILL) -
- TOOK FIVE QUAIL EGGS

NOTE IN A CORN (HILL) -
- TOOK FIVE QUAIL EGGS

NOTE IN A CORN (HILL) -
- TOOK FIVE QUAIL EGGS

NOTE IN A CORN (HILL) -
- TOOK FIVE QUAIL EGGS

NESTING OF EASTERN BLUEBIRD AS OBSERVED -

NEST BUILDING - 3 TO 7 DAYS

NO. EGGS - RANGE FROM 3 TO 5

INCUBATION PERIOD - 14 TO 18 DAYS

NEST LIFE - 15 TO 30 DAYS

ONLY FEMALE INCUBATES -

MALE ASSISTS IN FEEDING, HOWEVER -

PREFER LOW ELEVATIONS FOR NEST (4' TO 15')

NESTING MATERIALS - USUALLY GRASSES

✓
TOOK SIX HOUSE WREN EGGS FROM
A NEST IN A MAIL BOX APR. 7, '35 (FRESH)

✓
TOOK FOUR STARLING EGGS FROM A NEST
3' IN A HOLE IN AN APPLE TREE - APR. 8, '35
INCUBATED $\frac{1}{8}$

✓
TOOK FIVE PHOEBE EGGS FROM A NEST
ON CLIFFS ABOVE JAMES R. - APR. 14, '35
FRESH - NR. 85'

✓
FOUND OVENBIRD'S NEST WITH FOUR
YOUNG (1 WK. OLD) AND ONE UNHATCHED EGG
IN WOODS ON SIDE OF HILL NEAR V.E.S.
APRIL 15, '35

✓
- TOOK FIVE STARLING EGGS FROM A
NEST IN A LOCUST (35') - FRESH - APR. 30, '35

✓
SAW WOOD THRUSH'S NEST WITH FOUR
EGGS - MAY 1, '35. ALSO SONG SPARROW
AND CHIPPING SPARROWS NESTS WITH
FIVE AND ONE EGGS RESPECTIVELY.

1. *Chrysomelidae*
 2. *Curculionidae*
 3. *Chrysomelidae*
 4. *Curculionidae*
 5. *Chrysomelidae*
 6. *Curculionidae*
 7. *Chrysomelidae*
 8. *Curculionidae*
 9. *Chrysomelidae*
 10. *Curculionidae*
 11. *Chrysomelidae*
 12. *Curculionidae*
 13. *Chrysomelidae*
 14. *Curculionidae*
 15. *Chrysomelidae*
 16. *Curculionidae*
 17. *Chrysomelidae*
 18. *Curculionidae*
 19. *Chrysomelidae*
 20. *Curculionidae*
 21. *Chrysomelidae*
 22. *Curculionidae*
 23. *Chrysomelidae*
 24. *Curculionidae*
 25. *Chrysomelidae*
 26. *Curculionidae*
 27. *Chrysomelidae*
 28. *Curculionidae*
 29. *Chrysomelidae*
 30. *Curculionidae*
 31. *Chrysomelidae*
 32. *Curculionidae*
 33. *Chrysomelidae*
 34. *Curculionidae*
 35. *Chrysomelidae*
 36. *Curculionidae*
 37. *Chrysomelidae*
 38. *Curculionidae*
 39. *Chrysomelidae*
 40. *Curculionidae*
 41. *Chrysomelidae*
 42. *Curculionidae*
 43. *Chrysomelidae*
 44. *Curculionidae*
 45. *Chrysomelidae*
 46. *Curculionidae*
 47. *Chrysomelidae*
 48. *Curculionidae*
 49. *Chrysomelidae*
 50. *Curculionidae*
 51. *Chrysomelidae*
 52. *Curculionidae*
 53. *Chrysomelidae*
 54. *Curculionidae*
 55. *Chrysomelidae*
 56. *Curculionidae*
 57. *Chrysomelidae*
 58. *Curculionidae*
 59. *Chrysomelidae*
 60. *Curculionidae*
 61. *Chrysomelidae*
 62. *Curculionidae*
 63. *Chrysomelidae*
 64. *Curculionidae*
 65. *Chrysomelidae*
 66. *Curculionidae*
 67. *Chrysomelidae*
 68. *Curculionidae*
 69. *Chrysomelidae*
 70. *Curculionidae*
 71. *Chrysomelidae*
 72. *Curculionidae*
 73. *Chrysomelidae*
 74. *Curculionidae*
 75. *Chrysomelidae*
 76. *Curculionidae*
 77. *Chrysomelidae*
 78. *Curculionidae*
 79. *Chrysomelidae*
 80. *Curculionidae*
 81. *Chrysomelidae*
 82. *Curculionidae*
 83. *Chrysomelidae*
 84. *Curculionidae*
 85. *Chrysomelidae*
 86. *Curculionidae*
 87. *Chrysomelidae*
 88. *Curculionidae*
 89. *Chrysomelidae*
 90. *Curculionidae*
 91. *Chrysomelidae*
 92. *Curculionidae*
 93. *Chrysomelidae*
 94. *Curculionidae*
 95. *Chrysomelidae*
 96. *Curculionidae*
 97. *Chrysomelidae*
 98. *Curculionidae*
 99. *Chrysomelidae*
 100. *Curculionidae*

Woodcock
 Owl, Barn
 Owl, Long Eared
 Owl, Barred
 Crow, American
 1-12
 WMA

Crow, American
 1-12
 WMA

Woodcock
 Owl, Barn
 Owl, Long Eared
 Owl, Barred
 Crow, American
 1-12
 WMA

Crow, American
 1-12
 WMA

Woodcock
 Owl, Barn
 Owl, Long Eared
 Owl, Barred
 Crow, American
 1-12
 WMA

Crow, American
 1-12
 WMA

Woodcock
 Owl, Barn
 Owl, Long Eared
 Owl, Barred
 Crow, American
 1-12
 WMA

MARCH 15-30

Woodcock
 Owl, Barn
 Owl, Long Eared
 Owl, Barred

Crow, American

APRIL 1-15

* Duck, Wood
 Hawk, Red Tailed
 Hawk, Red Should.
 Owl, Barn
 Woodpecker, Hairy
 Titmouse, Tufted

Owl, Long Eared
 Owl, Barred
 Crow, Amer.
 Owl, Screech
 Woodpecker, Downy
 Chickadee, Carol.

15-30

Heron, Gt. Blue
 Heron, Green
 Heron, Blk. C.N.
 Rail, Clapper
 Shrike, Migrant
 Wren, Carolina
 Towhee

Woodpecker, Hairy
 Chickadee, Downy
 Sapsucker, V.B.
 Duck, Black
 Duck, Wood
 Nuthatch, W.Br.

MAY 1-15

Duck, Pintail
 * Bittern, American
 Heron, Little Blue
 " , Gt. Blue
 " , Green
 " , Bl. C.N.
 Grouse, Ruffed

* Hawk, Marsh
 Hawk, S.S.
 Hawk, Br. Winged
 Owl, Sh. Eared
 * Crow, Fish
 Martin, Purple
 Creeper, Brown

ORTHOPTERA-

Grasshoppers - ACRIDIDAE - 2 { NYMPH 4
ADULT

TETTIGONIIDAE -

KATYDID

CRICKETS -

COCKROACH -

HEMIPTERA -

CICADAS -

- BIRD'S EGGS -

Carbonate of lime - chief const.
phosphate of lime & magnesia
3 layers traversed by vertical
lines to snake pores.
Third layer a ^(ostrich) glaze or
porcelain or chalky (cuckoo)
some are translucent while
others very thick.

Eggs of ground laying birds are
usually pear shaped. While
eggs of others are more
or less shaped like a
hen's egg. Practically all
eggs laid in holes are white
& almost perfect spheres.
Some like chimney swift
are long & pointed on both
ends.

Usually 7 pigments constitute
colors in eggs. - brown^{1st} - red,
second -; deep blue - 3rd; light blue
yellow & rufous, 4th & 5th
combining in fifth to give a
green; brick red 6th; 7th is
nameless but gives a brown
tint when mixed with others.
Origin undetermined but
thought to arise from the
hemoglobin of blood.

Most eggs have surface
marked by spots or lines

or a combination of both: the pigment is deposited on the egg as it passes down the lower portion of the oviduct. The egg is believed to turn on its long axis as shown by spiral nature of spots & lines. Many eggs seem to have fainter markings than the outside markings, these are in the second layer of the egg; such eggs are said to be double spotted - as rails & plovers.

Among birds of prey the coloring of each egg seems to be more uniform & deeply colored. When but two eggs are laid most of dye is deposited on first & sometimes on second.



Coloring bears no relation to color of bird, ^{but more} ~~more~~ to its nature, of environment & hole nesting birds.

Ground " " "

Most birds build nests. Size of eggs depend mostly on condition of young at hatching. No. laid cannot ^{exceed} ~~exceed~~ no. that bird can cover. Invid zone

^{Immature}
Lays more & smaller eggs than
a fully developed ~~new~~ birds
megapodes (wild turkeys
of Austral.) ~~to~~ deposit eggs in
decaying veg. matter or in sand
around hot springs. All other
birds incubate requiring 2
~~weeks~~ to 6 weeks. Young
born practically helpless and
contained under the term
altricial requiring about 2 weeks
young born able to shift for
themselves known as praecocial
& require about 4 to 6 weeks to
hatch.

Composition of "Before Club"
Delivered before Club
The Science of May 5, 1935
The A. C. S.



The Efficiency Line



of Commercial Stationery